



INSTRUCTION FOR USE AND STORAGE OF THE STIMULATOR BIOWERM TECHNO

Scope of use, method, timing and doses of the BIOWERM TECHNO stimulator

BIOWERM TECHNO is an organic-mineral growth stimulator produced from vermicompost. It is intended for use in field crops and vegetable cultivation.

Dosage sizes and application timings:

Crop	Dose	Timing of application	Recommended amount water
Corn	2 l/ha	phase 2-3 leaves; phase 6- 8 leaves	200 L
Oil crops e.g.: Canola, Sunflower, Flax, Nigella	2 l/ha	Canola: <u>autumn</u> : phase 4-5 leaves; <u>spring</u> : after the start of vegetation; budding phase. Sunflower: phase of beginning of inflorescence development	200 L
Cereals e.g.: wheat, barley, oats, rye	2 l/ha	shoot phase; flag leaf phase	200 L
Vegetables , e.g.: zucchini, carrot, onion, radish, eggplant, watermelon, potatoes, beet, sugar beet, cabbage, broccoli, dill, parsley;	2 l/ha	Use for the <u>first time</u> : when the plant has 3-5 leaves; Use it for the second time after 7-15 days.	200 L
Fruit trees , e.g.: apple trees, plums, cherries, peaches, mulberry,	0,5-1 l/ha	<u>first time</u> : before flowering, <u>second time</u> immediately after flowering, <u>third time</u> : from as soon as the crop sets fruit	200 L
Berries , e.g.: strawberries, blueberries, currants	2 l/ha	<u>First time</u> : after vegetation; <u>Second time</u> : 5-7 days after first application	200 L



Method of preparing the working solution

Read the label before use.

Shake the bottle well before use.

Fill the working tank halfway with water and start mixing. Add the required amount of BLOWERM TECHNO to the sprayer tank, mix until dissolved or dispersed, and add the remaining amount of water. Make sure mixing is maintained during spraying.

Rinse the emptied container three times with water, pour the rinsings into the sprayer tank with the working solution. Wash the sprayer after use.

Notes

- Do not use on frozen, waterlogged, saturated, snow-covered soils or during rainfall. Soils that thaw at least on the surface during the day are not considered frozen.
- Do not use together with plant protection products.
- Do not exceed the recommended doses.

Storage

The product should be stored:

- in a closed, original container,
- at a temperature from +5°C to +15°C,
- away from sources of heat, oxidizing agents, (strong) acids
- out of reach of unauthorized persons, children, and animals.

Do not store with food, beverages, feed, or water.

Precautions for safe handling of the product

During When working with the stimulator, use protective gloves and eye protection. Avoid contact with skin, eyes, and clothing. In case of eye contact, rinse with plenty of water and seek medical advice. In case of skin contact, wash with plenty of water and soap. If irritation occurs, contact a doctor.

After work, wash hands with soap and water. Remove and wash contaminated clothing before reuse.

Akceptuję treść instrukcji stosowania
w zakresie upraw polowych
20.08.2021

Damian Wach
dr Damian Wach

Instytut Uprawy Nawożenia i Zołozenia
Państwowy Instytut Badawczy
ZAKŁAD ŻYWIENIA ROŚLIN
I NAWOZENA
24-100 Puławy, ul. Czeretoryskich 8
Tel. (81) 47 86 830